

# Spatially structured simulated community data

## Source of data

This dataset was created using the library `neutral.vp`, written by [Tyler Smith](#). This library implements the original algorithm for simulation of spatially structured multivariate data, developed by Graham Bell (2000, 2003 and 2005) for simulation of communities assembled by neutral processes. Smith & Lundholm (2010) extended Bell's algorithm by adding niche-based processes (see their paper for details).

The library `neutral.vp` was published as an [on-line appendix](#) of Smith & Lundholm (2010); published version contains [library source](#), which needs to be compiled<sup>1)</sup> (I have done this for the purpose of this study materials - see the link below).

## Description of the dataset


Simulated data represent an artificial landscape with 50 simulated species. The platform for landscape simulation was the topographical map of SW Moravia in Czech Republic (which is cut through by several deep river valleys in NW to SE direction). The artificial landscape is of size 100 x 100 pixels, each pixel representing one habitat with specific value of environmental variable (here elevation, rescaled to 1-10 scale). Each habitat (pixel) has an upper carrying capacity of 500 individuals, and each habitat is internally homogeneous. At the beginning of the simulation, 500 individuals (the same number for each of 50 species) are assigned to each habitat. Each species has randomly assigned ecological optima (on the scale 1-10), which influences the species survival at the habitat: the more similar is the species optima to the value of the habitat, the higher is the probability that species will produce an offspring (higher *birth rate*) and lower probability that the species will become extinct at the habitat (lower *death rate*). The offspring spreads to surrounding habitats by *random walk*; the higher is the migration speed set up in the model, the farther the offspring will get.

The model has two turning knobs - the first knob can regulate the importance of the environment (*niche-assembly*, i.e., how much is the survival of the species influenced by the difference between species ecological niche and habitat actual ecological conditions), and the second knob modifies the migration speed (*dispersal limitation*, i.e., how far the offspring can migrate).

The landscape is sampled in a regular fashion (only 144 habitats, separated by lag distance of 8 pixels, are sampled). The species data were created using the library `neutral.vp`, the script can be found [here](#).

## Environmental variables

The information about environmental conditions of individual habitats (pixels) is represented by elevation in topographical map of SW Moravia (100 x 100 pixels).



Topographical map of SW Moravia

## Data for download

R library [neutral.vp\\_1.0-0.zip](#) (author: Tyler Smith, published as an appendix to Smith & Lundholm 2010, current version compiled in R 3.1.0; x86\_64-w64-mingw32)<sup>2)</sup>. The library requires installation of additional libraries: [vegan](#), [permute](#) and [packfor](#)<sup>3)</sup>.

## Skript for direct import of data to R

```
low.disp.spe <- read.delim
('http://www.sci.muni.cz/botany/zeleny/wiki/anadat-r/lib/exe/fetch.php?media
=data:low-disp-spe.txt')
high.disp.spe <- read.delim
('http://www.sci.muni.cz/botany/zeleny/wiki/anadat-r/lib/exe/fetch.php?media
=data:high-disp-spe.txt')
env <- read.delim
('http://www.sci.muni.cz/botany/zeleny/wiki/anadat-r/lib/exe/fetch.php?media
=data:env-spatial-model.txt')
coord <- read.delim
('http://www.sci.muni.cz/botany/zeleny/wiki/anadat-r/lib/exe/fetch.php?media
=data:coord-spatial-model.txt')
```

## References

- Bell G. (2000): The distribution of abundance in neutral communities. *Am. Nat.* 155: 606–617.
- Bell G. (2003) The interpretation of biological surveys. *Proc. R. Soc. B* 270: 2531–2542.
- Bell G. (2005) The co-distribution of species in relation to the neutral theory of community ecology. *Ecology* 86: 1757-1770.
- Smith T.W. & Lundholm J.T. (2010): Variation partitioning as a tool to distinguish between niche and neutral processes. *Ecography* 33: 648-655 [pdf](#)

1)

To compile the library from source, you need a bunch of other software - so called Rtools, see [Windows toolset appendix](#).

2)

Previous version of compiled package, compiled in R 2.15.0, can be found [here](#)

3)

packfor should be installed from [R-forge](#) using the function `install.packages("packfor", repos="http://R-Forge.R-project.org")`. However, it works only for the latest version of R; if you need to install the library to an older R version, you need to compile it by your self from the source (the file .tar.gz).

From:

<https://anadat-r.davidzeleny.net/> - **Analysis of community ecology data in R**

Permanent link:

<https://anadat-r.davidzeleny.net/doku.php/en:data:simul-spatial?rev=1406209170>

Last update: **2017/10/11 12:36**